

Ch. 3: Descriptive Statistics

$$\bar{x} = \frac{\sum x}{n} \text{ Mean}$$

$$\bar{x} = \frac{\sum f \cdot x}{\sum f} \text{ Mean (frequency table)}$$

$$s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}} \text{ Standard deviation}$$

$$s = \sqrt{\frac{n(\sum x^2) - (\sum x)^2}{n(n - 1)}} \text{ Standard deviation (shortcut)}$$

$$s = \sqrt{\frac{n[\sum (f \cdot x^2)] - [\sum (f \cdot x)]^2}{n(n - 1)}} \text{ Standard deviation (frequency table)}$$

variance = s^2

Ch. 4: Probability

$P(A \text{ or } B) = P(A) + P(B)$ if A, B are mutually exclusive

$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$ if A, B are not mutually exclusive

$P(A \text{ and } B) = P(A) \cdot P(B)$ if A, B are independent

$P(A \text{ and } B) = P(A) \cdot P(B|A)$ if A, B are dependent

$P(\bar{A}) = 1 - P(A)$ Rule of complements

$nPr = \frac{n!}{(n - r)!}$ Permutations (no elements alike)

$\frac{n!}{n_1! n_2! \dots n_k!}$ Permutations (n_1 alike, ...)

$nCr = \frac{n!}{(n - r)! r!}$ Combinations

Ch. 5: Probability Distributions

$\mu = \sum x \cdot P(x)$ Mean (prob. dist.)

$\sigma = \sqrt{\sum [x^2 \cdot P(x)] - \mu^2}$ Standard deviation (prob. dist.)

$P(x) = \frac{n!}{(n - x)! x!} \cdot p^x \cdot q^{n-x}$ Binomial probability

$\mu = n \cdot p$ Mean (binomial)

$\sigma^2 = n \cdot p \cdot q$ Variance (binomial)

$\sigma = \sqrt{n \cdot p \cdot q}$ Standard deviation (binomial)

Ch. 6: Normal Distribution

$z = \frac{x - \bar{x}}{s}$ or $\frac{x - \mu}{\sigma}$ Standard score

$\mu_2 = \mu$ Central limit theorem

$\sigma_2 = \frac{\sigma}{\sqrt{n}}$ Central limit theorem (Standard error)

Ch. 7: Confidence Intervals (one population)

$\hat{p} - E < p < \hat{p} + E$ Proportion

where $E = z_{\alpha/2} \sqrt{\frac{\hat{p}\hat{q}}{n}}$

$\bar{x} - E < \mu < \bar{x} + E$ Mean

where $E = z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ (σ known)

or $E = t_{\alpha/2} \frac{s}{\sqrt{n}}$ (σ unknown)

$\frac{(n - 1)s^2}{\chi^2_k} < \sigma^2 < \frac{(n - 1)s^2}{\chi^2_l}$ Variance

Ch. 7: Sample Size Determination

$n = \frac{[z_{\alpha/2}]^2 \cdot 0.25}{E^2}$ Proportion

$n = \frac{[z_{\alpha/2}]^2 \hat{p}\hat{q}}{E^2}$ Proportion (\hat{p} and \hat{q} are known)

$n = \left[\frac{z_{\alpha/2} \sigma}{E} \right]^2$ Mean

Ch. 9: Confidence Intervals (two populations)

$(\hat{p}_1 - \hat{p}_2) - E < (p_1 - p_2) < (\hat{p}_1 - \hat{p}_2) + E$

where $E = z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}$

$(\bar{x}_1 - \bar{x}_2) - E < (\mu_1 - \mu_2) < (\bar{x}_1 - \bar{x}_2) + E$ (Indep.)

where $E = t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$ ($df = \text{smaller of } n_1 - 1, n_2 - 1$)

(σ_1 and σ_2 unknown and not assumed equal)

$E = t_{\alpha/2} \sqrt{\frac{s_p^2}{n_1} + \frac{s_p^2}{n_2}}$ ($df = n_1 + n_2 - 2$)

$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{(n_1 - 1) + (n_2 - 1)}$

(σ_1 and σ_2 unknown but assumed equal)

$E = z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$

(σ_1, σ_2 known)

$\bar{d} - E < \mu_d < \bar{d} + E$ (Matched pairs)

where $E = t_{\alpha/2} \frac{s_d}{\sqrt{n}}$ ($df = n - 1$)

Ch. 8: Test Statistics (one population)

$z = \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}}$ Proportion—one population

$z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$ Mean—one population (σ known)

$t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$ Mean—one population (σ unknown)

$\chi^2 = \frac{(n - 1)s^2}{\sigma^2}$ Standard deviation or variance—one population

Ch. 9: Test Statistics (two populations)

$z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{\hat{p}\hat{q}}{n_1} + \frac{\hat{p}\hat{q}}{n_2}}}$ Two proportions

where $\hat{p} = \frac{x_1 + x_2}{n_1 + n_2}$

$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$ $df = \text{smaller of } n_1 - 1, n_2 - 1$

Two means—dependent; σ_1 and σ_2 unknown, and not assumed equal.

$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_p^2}{n_1} + \frac{s_p^2}{n_2}}}$ ($df = n_1 + n_2 - 2$)

Two means—dependent; σ_1 and σ_2 unknown, but assumed equal.

$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$ Two means—dependent; σ_1, σ_2 known.

$t = \frac{\bar{d} - \mu_d}{s_d/\sqrt{n}}$ Two means—matched pairs ($df = n - 1$)

Ch. 11: Goodness-of-Fit and Contingency Tables

$\chi^2 = \sum \frac{(O - E)^2}{E}$ Goodness-of-fit ($df = k - 1$)

$\chi^2 = \sum \frac{(O - E)^2}{E}$ Contingency table ($df = (r - 1)(c - 1)$)

where $E = \frac{(\text{row total})(\text{column total})}{(\text{grand total})}$

Ch. 10: Linear Correlation/Regression

Correlation $r = \frac{n\sum xy - (\sum x)(\sum y)}{\sqrt{[n\sum x^2 - (\sum x)^2][n\sum y^2 - (\sum y)^2]}}$

or $r = \frac{\sum (z_x z_y)}{n - 1}$ where $z_x = z$ score for x , $z_y = z$ score for y

Slope: $b_1 = \frac{n\sum xy - (\sum x)(\sum y)}{n\sum x^2 - (\sum x)^2}$

or $b_1 = r \frac{s_y}{s_x}$

y-Intercept: $b_0 = \bar{y} - b_1 \bar{x}$ or $b_0 = \frac{(\sum y)(\sum x^2) - (\sum x)(\sum xy)}{n(\sum x^2) - (\sum x)^2}$

$\hat{y} = b_0 + b_1 x$ Estimated eq. of regression line

$r^2 = \frac{\text{explained variation}}{\text{total variation}}$

$t_r = \frac{\sqrt{\sum (y - \hat{y})^2}}{n - 2}$ or $\sqrt{\frac{\sum y^2 - b_0 \sum y - b_1 \sum xy}{n - 2}}$

$\hat{y} - E < y < \hat{y} + E$ Prediction interval

where $E = t_{\alpha/2} \sqrt{1 + \frac{1}{n} + \frac{n(x_0 - \bar{x})^2}{n\sum x^2 - (\sum x)^2}}$

$r_s = 1 - \frac{6\sum d^2}{n(n^2 - 1)}$ Rank correlation

(critical value for $n > 30$: $\frac{\pm z}{\sqrt{n - 1}}$)

Ch. 11: One-Way Analysis of Variance

Procedure for testing $H_0: \mu_1 = \mu_2 = \mu_3 = \dots$

- Use software or calculator to obtain results.
- Identify the P -value.
- Form conclusion:
 - If P -value $\leq \alpha$, reject the null hypothesis of equal means.
 - If P -value $> \alpha$, fail to reject the null hypothesis of equal means.

TABLE A-3 t Distribution: Critical t Values

Degrees of Freedom	Area in One Tail				
	0.005	0.01	0.025	0.05	0.10
1	63.657	31.821	12.706	6.314	3.078
2	9.925	6.965	4.303	2.920	1.886
3	5.841	4.541	3.182	2.353	1.638
4	4.604	3.747	2.776	2.132	1.533
5	4.032	3.365	2.571	2.015	1.476
6	3.707	3.143	2.447	1.943	1.440
7	3.499	2.998	2.365	1.895	1.415
8	3.355	2.896	2.306	1.860	1.397
9	3.250	2.821	2.262	1.833	1.383
10	3.169	2.764	2.228	1.812	1.372
11	3.106	2.718	2.201	1.796	1.363
12	3.055	2.681	2.179	1.782	1.356
13	3.012	2.650	2.160	1.771	1.350
14	2.977	2.624	2.145	1.761	1.345
15	2.947	2.602	2.131	1.753	1.341
16	2.921	2.583	2.120	1.746	1.337
17	2.898	2.567	2.110	1.740	1.333
18	2.878	2.552	2.101	1.734	1.330
19	2.861	2.539	2.093	1.729	1.328
20	2.845	2.528	2.086	1.725	1.325
21	2.831	2.518	2.080	1.721	1.323
22	2.819	2.508	2.074	1.717	1.321
23	2.807	2.500	2.069	1.714	1.319
24	2.797	2.492	2.064	1.711	1.318
25	2.787	2.485	2.060	1.708	1.316
26	2.779	2.479	2.056	1.706	1.315
27	2.771	2.473	2.052	1.703	1.314
28	2.763	2.467	2.048	1.701	1.313
29	2.756	2.462	2.045	1.699	1.311
30	2.750	2.457	2.042	1.697	1.310
31	2.744	2.453	2.040	1.696	1.309
32	2.738	2.449	2.037	1.694	1.309
33	2.733	2.445	2.035	1.692	1.308
34	2.728	2.441	2.032	1.691	1.307
35	2.724	2.438	2.030	1.690	1.306
36	2.719	2.434	2.028	1.688	1.306
37	2.715	2.431	2.026	1.687	1.305
38	2.712	2.429	2.024	1.686	1.304
39	2.708	2.426	2.023	1.685	1.304
40	2.704	2.423	2.021	1.684	1.303
45	2.690	2.412	2.014	1.679	1.301
50	2.678	2.403	2.009	1.676	1.299
60	2.660	2.390	2.000	1.671	1.296
70	2.648	2.381	1.994	1.667	1.294
80	2.639	2.374	1.990	1.664	1.292
90	2.632	2.368	1.987	1.662	1.291
100	2.626	2.364	1.984	1.660	1.290
200	2.601	2.345	1.972	1.653	1.286
300	2.592	2.339	1.968	1.650	1.284
400	2.588	2.336	1.966	1.649	1.284
500	2.586	2.334	1.965	1.648	1.283
1000	2.581	2.330	1.962	1.646	1.282
2000	2.578	2.328	1.961	1.646	1.282
Large	2.576	2.326	1.960	1.645	1.282

TABLE A-5 Critical Values of the Pearson Correlation Coefficient r

n	$\alpha = 0.05$	$\alpha = 0.01$
4	.950	.990
5	.878	.959
6	.811	.917
7	.754	.875
8	.707	.834
9	.666	.798
10	.632	.765
11	.602	.735
12	.576	.708
13	.553	.684
14	.532	.661
15	.514	.641
16	.497	.623
17	.482	.606
18	.468	.590
19	.456	.575
20	.444	.561
25	.396	.505
30	.361	.463
35	.335	.430
40	.312	.402
45	.294	.378
50	.279	.361
60	.254	.330
70	.236	.305
80	.220	.286
90	.207	.269
100	.196	.256

NOTE: To test $H_0: \rho = 0$ against $H_1: \rho \neq 0$, reject H_0 if the absolute value of r is greater than the critical value in the table.

NEGATIVE z Scores





TABLE A-2 Standard Normal (z) Distribution: Cumulative Area from the LEFT

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.50 and lower	.0001									
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0006	.0006	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0010	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0014	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0020	.0019	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

NOTE: For values of z below -3.49, use 0.0001 for the area.

*Use these common values that result from interpolation:

z score	Area
-1.645	0.0500
-2.575	0.0050



POSITIVE z Scores

TABLE A-2 (continued) Cumulative Area from the LEFT.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9995	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998
3.50 and up	.9999									

NOTE: For values of z above 3.49, use 0.9999 for the area.

*Use these common values that result from interpolation:

z score	Area
1.645	0.9500
2.575	0.9950

Common Critical Values

Confidence Level	Critical Value
0.90	1.645
0.95	1.96
0.99	2.575

TABLE A-4 Chi-Square (χ^2) Distribution
Area to the Right of the Critical Value

Degrees of Freedom	Area to the Right of the Critical Value									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	0.0001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
2	0.0100	0.0200	0.0501	0.1003	0.2101	0.4457	0.7173	1.0545	1.3858	1.8477
3	0.0778	0.1150	0.2167	0.3527	0.5844	1.0085	1.4959	2.3658	3.2187	4.1017
4	0.2072	0.2978	0.4844	0.7173	1.0641	1.4860	2.2041	3.3571	4.6079	5.7928
5	0.4120	0.5544	0.8311	1.1455	1.6013	2.2041	3.3571	4.6079	6.2514	7.8798
6	0.6758	0.8721	1.2372	1.6354	2.2041	3.3571	4.6079	6.2514	8.0538	9.8878
7	0.9893	1.2398	1.6900	2.1673	2.8331	3.8282	5.0413	6.3441	8.0335	9.8906
8	1.3444	1.6466	2.1800	2.7333	3.4900	4.6079	6.2514	8.0335	9.8906	12.0170
9	1.7350	2.0888	2.7000	3.3355	4.1682	5.4078	7.1721	9.0032	11.3445	13.5810
10	2.1561	2.5588	3.2470	3.9400	4.8652	6.1799	8.1912	10.4845	12.9360	15.4543
11	2.6033	3.0533	3.8166	4.5750	5.5780	7.1721	9.0032	11.3445	13.5810	16.5910
12	3.0704	3.5711	4.4004	5.2236	6.3044	8.0335	10.2157	12.5916	14.6976	17.7793
13	3.5652	4.1070	5.0099	5.8922	7.0420	9.2364	11.5780	14.1645	16.2139	19.0291
14	4.0750	4.6660	5.6299	6.5270	7.7900	10.0420	13.0009	15.7857	17.7600	20.3390
15	4.6011	5.2299	6.2620	7.2610	8.5410	11.1548	14.3380	17.5345	19.5912	21.7822
16	5.1420	5.8120	6.9008	7.9620	9.3120	12.4422	15.8585	19.5912	21.7822	23.3646
17	5.6875	6.4060	7.5640	8.6720	10.0850	13.8009	17.5345	21.7822	23.3646	25.0001
18	6.2850	7.0150	8.2310	9.3900	10.8650	15.3080	19.5912	23.3646	25.0001	26.7678
19	6.8440	7.6330	8.9070	10.1170	11.6510	16.9190	21.7822	25.0001	26.7678	28.5768
20	7.4340	8.2660	9.5910	10.8510	12.4420	18.5990	24.4330	28.5768	28.5768	30.4190
21	8.0340	8.8970	10.2830	11.5810	13.2400	20.3390	26.7678	30.4190	30.4190	32.2760
22	8.6430	9.5420	10.9820	12.3380	14.0420	22.1640	29.1910	32.2760	32.2760	34.1480
23	9.2600	10.1960	11.6890	13.0910	14.8480	24.0000	31.5260	34.1480	34.1480	36.0350
24	9.8860	10.8560	12.4010	13.8480	15.6590	25.7900	33.6780	36.0350	36.0350	37.9360
25	10.5200	11.5240	13.1200	14.6110	16.4730	27.5620	35.6520	37.9360	37.9360	39.8510
26	11.1600	12.1980	13.8440	15.3790	17.2920	29.3170	37.6660	39.8510	39.8510	41.7800
27	11.8080	12.8790	14.5730	16.1510	18.1140	31.0540	39.8120	41.7800	41.7800	43.7230
28	12.4610	13.5650	15.3080	16.9280	18.9390	32.7730	41.9030	43.7230	43.7230	45.6800
29	13.121									